



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Physiological Concentrations of Amyloid Beta Regulate Recycling of Synaptic Vesicles via Alpha7 Acetylcholine Receptor and CDK5/Calcineurin Signaling

Citation for published version:

Lazarevic, V, Fienko, S, Andres-Alonso, M, Anni, D, Ivanova, D, Montenegro-Venegas, C, Gundelfinger, ED, Cousin, MA & Fejtova, A 2017, 'Physiological Concentrations of Amyloid Beta Regulate Recycling of Synaptic Vesicles via Alpha7 Acetylcholine Receptor and CDK5/Calcineurin Signaling', *Frontiers in Molecular Neuroscience*, vol. 10, 221. <https://doi.org/10.3389/fnmol.2017.00221>

Digital Object Identifier (DOI):

[10.3389/fnmol.2017.00221](https://doi.org/10.3389/fnmol.2017.00221)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Frontiers in Molecular Neuroscience

Publisher Rights Statement:

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Physiological Concentrations of Amyloid Beta Regulate Recycling of Synaptic Vesicles via Alpha7 Acetylcholine Receptor and CDK5/Calcineurin Signaling

Vesna Lazarevic^{1,2,3†}, Sandra Fieńko^{1‡}, Maria Andres-Alonso^{1‡}, Daniela Anni^{4‡}, Daniela Ivanova¹, Carolina Montenegro-Venegas², Eckart D. Gundelfinger^{2,3,5,6}, Michael A. Cousin⁷ and Anna Fejtova^{1,2,4,5*}

OPEN ACCESS

Edited by:

Ulrike C. Müller,
Heidelberg University, Germany

Reviewed by:

Andreas Vlachos,
Albert Ludwig University of Freiburg,
Germany
Gael Barthelet,
UMR5297 Institut Interdisciplinaire de
Neurosciences (IINS), France

*Correspondence:

Anna Fejtova
anna.fejtova@uk-erlangen.de

†Present address:

Vesna Lazarevic,
Department of Clinical Neuroscience,
Center for Molecular Medicine,
Karolinska Institutet, Stockholm,
Sweden

‡These authors have contributed
equally to this work.

Received: 30 March 2017

Accepted: 26 June 2017

Published: 21 July 2017

Citation:

Lazarevic V, Fieńko S, Andres-Alonso M, Anni D, Ivanova D, Montenegro-Venegas C, Gundelfinger ED, Cousin MA and Fejtova A (2017) Physiological Concentrations of Amyloid Beta Regulate Recycling of Synaptic Vesicles via Alpha7 Acetylcholine Receptor and CDK5/Calcineurin Signaling. *Front. Mol. Neurosci.* 10:221. doi: 10.3389/fnmol.2017.00221

¹RG Presynaptic Plasticity, Leibniz Institute for Neurobiology, Magdeburg, Germany, ²Department of Neurochemistry and Molecular Biology, Leibniz Institute for Neurobiology, Magdeburg, Germany, ³German Center for Neurodegenerative Diseases (DZNE), Magdeburg, Germany, ⁴Molecular Psychiatry, Department of Psychiatry and Psychotherapy, University Hospital, University of Erlangen-Nuremberg, Erlangen, Germany, ⁵Center for Behavioral Brain Sciences, Otto von Guericke University, Magdeburg, Germany, ⁶Medical Faculty, Otto von Guericke University, Magdeburg, Germany, ⁷Centre for Integrative Physiology, University of Edinburgh, Edinburgh, United Kingdom

Despite the central role of amyloid β (A β) peptide in the etiopathogenesis of Alzheimer's disease (AD), its physiological function in healthy brain is still debated. It is well established that elevated levels of A β induce synaptic depression and dismantling, connected with neurotoxicity and neuronal loss. Growing evidence suggests a positive regulatory effect of A β on synaptic function and cognition; however the exact cellular and molecular correlates are still unclear. In this work, we tested the effect of physiological concentrations of A β species of endogenous origin on neurotransmitter release in rat cortical and hippocampal neurons grown in dissociated cultures. Modulation of production and degradation of the endogenous A β species as well as applications of the synthetic rodent A β_{40} and A β_{42} affected efficacy of neurotransmitter release from individual presynapses. Low picomolar A β_{40} and A β_{42} increased, while A β depletion or application of low micromolar concentration decreased synaptic vesicle recycling, showing a hormetic effect of A β on neurotransmitter release. These A β -mediated modulations required functional alpha7 acetylcholine receptors as well as extracellular and intracellular calcium, involved regulation of CDK5 and calcineurin signaling and increased recycling of synaptic vesicles. These data indicate that A β regulates neurotransmitter release from presynapse and suggest that failure of the normal physiological function of A β in the fine-tuning of SV cycling could disrupt synaptic function and homeostasis, which would, eventually, lead to cognitive decline and neurodegeneration.

Keywords: amyloid beta, acetylcholine receptors, synaptic vesicle recycling, neurotransmitter release, CDK5, calcineurin

INTRODUCTION

Amyloid beta (A β) peptide arises by processing of the extracellular domain of the amyloid precursor protein (APP) mediated by the β - and γ -secretase proteolytic activity. A β is famous for its connection with Alzheimer's disease (AD), the most common form of neurodegeneration, characterized by progressive cognitive decline, memory impairment and formation of amyloid plaques in the brains of affected patients. A β is the main component of these plaques (Glenner and Wong, 1984). This fact together with a genetic link of early-onset familial forms of AD to mutations interfering with the proteolytic processing of APP (Goate et al., 1991) advocate the key role of extracellular A β in the pathogenesis of AD. However, the mechanisms by which dysregulated extracellular A β disrupts the brain function are still not completely understood. While there is no clear relationship between the number of senile plaques and disease severity, progressive synapse dismantling occurring before formation of any amyloid deposits emerged as the best correlate of the cognitive decline in AD patients and animal models (Terry et al., 1991; Mucke et al., 2000). As synapse loss in AD is preceded by defects in neuronal transmission and plasticity, it has been suggested that extracellular A β might be involved in the regulation of these processes (Chapman et al., 1999; Hsia et al., 1999; Walsh et al., 2002; Palop and Mucke, 2010). A large body of evidence supports an inhibitory effect of A β on synaptic function. Elevated (high nanomolar and low micromolar) extracellular A β reduces neurotransmission mainly by postsynaptic mechanisms including increased internalization or desensitization of postsynaptic glutamate receptors and downstream signaling (Walsh et al., 2002; Hsieh et al., 2006). Recently, it has also been demonstrated that high nanomolar A β affects endocytosis of presynaptic neurotransmitter vesicles indicating that the inhibitory effect of A β on membrane trafficking is not restricted only to the postsynaptic compartment (Park et al., 2013).

In healthy brain, A β is released into the extracellular space depending on neuronal activity (Kamenetz et al., 2003; Cirrito et al., 2005). Thus, due to the widely accepted inhibitory nature of A β on the neurotransmission, it has been proposed that endogenous A β functions as a negative modulator of synaptic strength in a physiological feed-back mechanism preventing over excitation of brain circuits. Dysregulation of this homeostatic mechanism would trigger synaptic destabilization eventually leading to the development of AD (Palop and Mucke, 2010). However, extracellular concentrations of A β in normal brain have been estimated to low picomolar levels, far lower than the concentrations used in most studies showing the A β -induced synaptic depression and neurotoxicity (Cirrito et al., 2003; Puzzo et al., 2011). Unexpectedly, several studies investigating the impact of physiological concentration of A β revealed a positive effect on neuroplasticity and learning (Puzzo et al., 2008, 2011; Garcia-Osta and Alberini, 2009). Hippocampal long-term potentiation (LTP) and learning were improved upon application of synthetic mouse or human A β_{42} in picomolar concentrations (Puzzo et al., 2008, 2011; Garcia-Osta and Alberini, 2009), whereas high nanomolar A β , in the

same experimental setting, led to well-established reduction of LTP suggesting a hormetic nature of A β on synaptic plasticity (Puzzo et al., 2008). Interestingly, these effects were sensitive to α -bungarotoxin, a selective antagonist of α 7 nicotinic acetylcholine receptor (α 7nAChR), and absent in α 7nAChR knockout mice, which implies that functional α 7nAChRs are required for A β_{42} -induced neuroplasticity (Puzzo et al., 2008, 2011). This is consistent with the reported high-affinity binding of A β to α 7nAChR (Wang et al., 2000a,b) and increased calcium influx through α 7nAChRs in isolated hippocampal synaptosomes upon application of picomolar A β_{42} (Dougherty et al., 2003). It has been proposed that the positive impact of A β_{42} on neurotransmission is mediated by potentiating neurotransmitter release from presynapse (Puzzo et al., 2008; Abramov et al., 2009). Abramov et al. (2009) convincingly demonstrated modulation of presynaptic release probability in rat and mice hippocampal neuronal cultures treated with thiorphan (Th), an inhibitor of the rate-limiting peptidase neprilysin involved in the extracellular clearance of A β species. Later on, the same laboratory described the modulation of presynaptic release probability in the same rodent cultures upon application of picomolar amounts of human A β_{1-40} peptide (Abramov et al., 2009; Fogel et al., 2014). In contrast to *ex vivo* electrophysiological experiments in hippocampal slices and behavioral analyses (Puzzo et al., 2008, 2011), both studies in cultured cells argued against the contribution of α 7nAChR to the effect of Th or A β_{1-40} on neurotransmission and proposed an alternative pathway involving APP homodimerization and signaling via heteromeric G $_{i/o}$ proteins (Fogel et al., 2014).

Thus, it is unclear, whether different species of endogenous A β peptides exert the same effect on presynapse, what is the contribution of α 7nAChRs, and what signaling connects putative A β receptors to the regulation of neurotransmitter release from presynapse. To address these questions, we tested systematically presynaptic effects of Th and rodent A β_{1-40} as well as A β_{1-42} in low to intermediate picomolar and low micromolar concentrations in cultured cortical neurons. To this end we visualized and quantified synaptic vesicle (SV) recycling within individual presynaptic boutons in living cells and investigated the contribution of α 7nAChRs and their downstream signaling to the A β -mediated regulation of presynaptic function. Our data have potential implications for the pathophysiology of AD. Since A β modulates neurotransmission at very low extracellular concentrations, this physiological function would be directly affected already upon minor changes in extracellular A β levels occurring in early phases of AD and thus might contribute to cognitive impairments far before formation of amyloid plaques.

MATERIALS AND METHODS

Antibodies

For immunocytochemical stainings (ICC) and for Western blots (WB) following primary antibodies were used from rabbit: anti-CDK5 (WB 1:1000, C-8 Santa Cruz), anti-homer1 (ICC 1:1000, Synaptic Systems), anti-VGLUT1 (ICC 1:1000, Synaptic Systems), anti-VGAT (ICC 1:1000, Synaptic Systems),

anti-VGAT luminal domain Oyster550-labeled (ICC 1:200, Synaptic Systems), from mouse: anti-synaptotagmin1 luminal domain Oyster550-labeled (ICC 1:250, Synaptic Systems), anti- β -tubulin isotype III (WB 1:2000, Sigma), anti-A β _{17–24} (4G8) (5 μ g/ml, Signet), and from guinea pig: anti-synaptophysin (ICC 1:1000, Synaptic Systems). For ICC Alexa Fluor 488- (1:2000), Cy3- (1:2000) and Cy5- (1:1000) fluorescently labeled secondary antibodies were purchased from Jackson ImmunoResearch. For WB secondary antibodies labeled with Alexa Fluor 680 (1:20,000, ThermoFisher Scientific/Molecular Probes) and IRDye 800CW (1:20,000, Rockland) were used.

Chemical Reagents

Thiorphan (Th), FK-506 monohydrate, TMB8 and Choline chloride were purchased from Sigma-Aldrich. β -Secretase inhibitor IV, InSolution γ -Secretase inhibitor L-685, 458, InSolution Roscovitine, α -Bungarotoxin and Bafilomycin A1 from Calbiochem. A β _{1–42} and A β _{1–40} peptides, D-(-)-2-Amino-5-phosphonopentanoic acid (APV), 6-Cyano-7-nitroquinoxaline-2,3-dione disodium (CNQX), BAPTA-AM and PNU 120596 from Tocris. A β was diluted according to the manufacturer's instruction. Th was diluted to 1 mM stock solution in artificial cerebrospinal fluid (ASCF) supplemented with 1 mM ascorbic acid to prevent Th oxidation (Iwata et al., 2001; Abramov et al., 2009). In all experiments control cells were treated with ascorbic acid in ASCF.

Animals

Breeding of animals and experiments using animal material were carried out in accordance with the European Communities Council Directive (2010/63/EU) and approved by the local animal care committees of Sachsen-Anhalt and Middle-Franconia/Germany.

Primary Neuronal Cultures

Primary cultures of cortical neurons were prepared as described previously (Lazarevic et al., 2011). In brief, rat embryos at day 18–19 after fertilization (E18–E19) were sacrificed by decapitation. The brains were removed and deprived of meninges. After treatment with 0.25% trypsin for 15 min and mechanical trituration cell suspension was plated in DMEM containing 10% fetal calf serum (FCS), 1 mM glutamine and antibiotics (100 U/ml penicillin, 100 μ g/ml streptomycin) onto poly-D-lysine coated glass coverslips (Sigma, 18 mm diameter). Twenty-four hours after plating, the medium was exchanged for Neurobasal medium supplemented with B27 (Life Technologies), antibiotics, and 0.8 mM glutamine. The cells were maintained in a humidified incubator with 5% CO₂. Primary hippocampal cultures were prepared according to a modified original protocol from Banker (1980) as described in Frischknecht et al. (2008). Briefly, rat embryos were sacrificed at E18–E19, brains were removed, hippocampi extracted and subjected to trypsin digestion and mechanical trituration. Thereafter, cells suspended in DMEM containing 10% FCS, 1 mM glutamine and penicillin/streptomycin were plated onto poly-L-lysine-coated glass coverslips. After 1 h, coverslips with primary hippocampal neurons were transferred into a Petri dish containing an

astrocytic monolayer in Neurobasal medium supplemented with B27, antibiotics and glutamine, as described before, and placed in a humidified incubator. Neurons cultured for 18–21 days *in vitro* (DIV) were used for all analyses. For immunocytochemistry and A β ₄₂ ELISA cells were plated on poly-D-lysine-coated glass coverslips at a density of 50,000 cells/12 mm coverslip in 24-well plates in 0.5 ml growth media. A β ₄₀ ELISA was done on cells plated at density 100,000 cells/18 mm coverslips in 12-well plates in 1 ml of growth media. For imaging experiments cells were plated on poly-L-lysine-coated glass coverslips at a density of 30,000 cells/18 mm coverslip in a 60-mm Petri dish. For biochemical experiments cells were plated in 6-well plates at a density of 300,000 cells/well.

ELISA Measurements of A β

The concentration of A β peptides in extracellular medium was assessed by sandwich ELISA using Human/Rat A β (42) high sensitive kit and Human/Rat A β (40) kit II purchased from Wako. The measurements were done according to manufacturer's protocol always using fresh medium.

Immunocytochemistry and Synaptotagmin1 Luminal Domain Antibody Uptake

Neurons were fixed with 4% paraformaldehyde, 4% sucrose in PBS pH 7.4, for 3 min at RT. Prior to immunostaining, the cells were blocked and permeabilized with PBS containing 10% FCS, 0.1% glycine and 0.3% TritonX-100 for 30 min. Subsequently, primary antibodies were applied overnight at 4°C. After three washing steps with PBS at RT coverslips were incubated with secondary antibodies for 1 h at RT. Both, primary and secondary antibodies were diluted in PBS containing 3% FCS. Coverslips were mounted on slides with Mowiol (Calbiochem) and kept at 4°C until microscopic analysis. Synaptotagmin1 luminal domain antibody uptake was done as described (Lazarevic et al., 2011). Cells were briefly washed with freshly prepared Tyrode's buffer (119 mM NaCl, 2.5 mM KCl, 2 mM CaCl₂, 2 mM MgCl₂, 30 mM glucose, 25 mM HEPES pH 7.4) and incubated with fluorescently-labeled syt1 antibody diluted in the same buffer, either for 20 min at 37°C to monitor network activity driven uptake, or for 4 min at RT in Tyrode's buffer containing 50 mM KCl and 71.5 mM NaCl to assess evoked syt1 Ab uptake. Thereafter, samples were fixed and stained. In each experiment, at least two coverslips per treatment were processed in parallel. The results are representative of 2–5 independent experiments.

Lentiviral Particles Production

The original mRFP-synaptophysin-pHluorin (syphHy) construct was obtained T. Oertner (Rose et al., 2013). cDNA of SyphHy was inserted into FUGW backbone vector by standard cloning. Lentiviral particles were generated in HEK293T cell line (ATTC, Manassas, VA, USA) using FUGW-based transfer, psPAX2 packaging and pVSVG pseudotyping vectors (Lois et al., 2002). HEK293T cells were grown in media containing 10% FCS to 60% confluence in 75 cm² flasks. The cells were transfected using the calcium phosphate method (Fejtova et al., 2009). Molar

ratio of FUGW: psPAX2: pVSVG was 2:1:1. Twenty-four hours later the content of FCS was reduced to 4%. After 48 h, the virus-containing media was collected and cleared from large cellular debris by centrifugation for 20 min at 2000 g. The supernatant was aliquoted and stored at -80°C until further use.

Synapto-pHluorin Imaging

SypHy was delivered to rat hippocampal neurons cultured for 4 DIV by the means of lentiviral infection and subjected to imaging at DIV17–18. Coverslips were incubated with A β_{1-42} or water as a control in a conditioned media for 1 h in the incubator. Coverslips were installed in an imaging chamber (Warner Instruments) and imaged at RT on an inverted microscope (Observer. D1; Zeiss) endowed with an EMCCD camera (Evolve 512; Delta Photometrics) controlled by MetaMorph Imaging (MDS Analytical Technologies). A 63 \times objective and GFP/mCherry single band excitors ET filter set (exciter 470/40, exciter 572/35, emitter 59022m, dichroic 59022BS) were used. Transduced neurons were identified by RFP expression. Neurons were stimulated in the presence of bafilomycin A1 (1 μM) to prevent vesicle reacidification and APV (50 μM) and CNQX (10 μM) to block recurrent network activity with 40 AP at 20 Hz, followed by 2 min recovery period. Afterwards, a stimulation with 900 AP at 20 Hz was delivered and a pulse of 60 mM NH $_4\text{Cl}$ -containing solution applied (Burrone et al., 2006). Electrical stimulation was delivered by a S48 stimulator unit (GRASS Technologies). A stream of images was acquired at 10 Hz and 5 s of the baseline was recorded before each stimulation, followed by imaging of the recovery phase for another 60 s. Synaptic boutons responsive to stimulation were selected by subtracting the first 10 frames of the baseline (before stimulation) from the 10 frames directly after the stimulus. Only neurons showing $\leq 20\%$ increase in the fluorescence after NH $_4\text{Cl}$ application were considered as viable and metabolically active and included for the analysis. The mean IF intensities were measured in the circular regions of interest (ROIs with a diameter of 8×8 pixel) placed over each responding synapse using Time Series Analyzer V2.0 plugin in ImageJ and plotted after bleaching correction using GraphPad Software. The relative sizes of the RRP and the RP were expressed as fractions of the total sypHy-expressing pool detected after addition of NH $_4\text{Cl}$. RP was quantified by averaging the mean of 50 values per each cell (representative of the frames 390–440 corresponding to time points 39–44 s on the XY graph). The results are representative of 3–5 independent experiments.

Image Acquisition and Analysis

Images were acquired with Zeiss Axio Imager A2 microscope with Cool Snap EZ camera (Visitron Systems) and MetaMorph Imaging software (MDS Analytical Technologies). For each pair of coverslips (treated vs. control) the same exposure time was taken. Per each experimental condition two coverslips were individually treated and processed. Images were captured from at least 3–5 visual fields (= cells) per coverslip and further analyzed using NIH ImageJ and OpenView software (Tsuriel et al., 2006). Upon appropriate background subtraction, immunoreactive puncta were counted along the 20 μm of proximal ($\geq 10 \mu\text{m}$ and $\leq 50 \mu\text{m}$ distance from the cell body) or distal dendrite ($\geq 50 \mu\text{m}$

distance from the cell body). The synaptic immunofluorescence intensities (IF) were assessed in a region of interest (ROI) set by the mask in the channel for synaptophysin (sph), which was used as synaptic marker. The mask was created semiautomatically using OpenView software.

Quantitative Western Blot

Control and treated 3 weeks old cortical neurons were washed with ice cold PBS and lysed in lysis buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA) supplemented with Complete protease inhibitors (Roche), phosphatase inhibitor cocktail PhosStop (Roche) and Calpain inhibitor PD150606 (Tocris). Precleared cell lysates were mixed with SDS loading buffer. Equal amount of proteins was loaded on SDS-PAGE and electrotransferred to Millipore Immobilon-FL PVDF membranes. Membranes were incubated over night at 4°C with primary antibodies and 1 h at RT with fluorescently labeled secondary antibody. Immunodetection and quantification was carried out using Odyssey Infrared Imagine System and Odyssey software v2.1 (LI-COR). After appropriate background subtraction all values were normalized using β III-tubulin as a loading control.

CDK5 Immunoprecipitation and Activity Assay

The kinase assay was performed as described in Crews et al. (2011). Briefly, 3 weeks old cortical neurons were washed with ice cold PBS and lysed in CDK5 IP buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA) supplemented with Complete protease inhibitors, phosphatase inhibitor cocktail PhosStop and Calpain inhibitor PD150606. Precleared cell lysates were subjected to immunoprecipitation using GammaBind Plus Sepharose beads (GE Healthcare) coupled with rabbit polyclonal antibody against CDK5 (C-8; Santa Cruz Biotechnology). After 3 h incubation at 4°C , immunoprecipitates were washed three times with CDK5 IP buffer and resuspended in 50 μl of CDK5 kinase buffer (25 mM Tris pH 7.5, 10 mM MgCl $_2$) in the presence of 90 μM ATP and 0.1 mM CDK5 substrate, Histone H1 (PKTPKKAKKL; sc-3066 Santa Cruz). Samples were then incubated for 30 min at 30°C and the reaction was stopped by adding 50 μl of Kinase glo plus reagent (Promega). Luminescent signal was measured on FLUOstar Omega microplate reader (BMG Labtech).

Calcineurin Activity Assay

Calcineurin activity was assessed using calcineurin cellular activity assay kit (Calbiochem, Cat. No. 207007) according to the manufacturer's instructions. In brief, 3 weeks old primary cortical neurons were lysed in the buffer supplied by the manufacturer (25 mM Tris-HCl pH 7.5, 0.5 mM dithiothreitol, 50 μM EDTA, 50 μM EGTA, 0.2% Nonidet P-40). Upon removal of free phosphate samples were incubated with calcineurin substrate, RII phosphopeptide (DLDVIPGRFDRRVpSVAAE), in the assay buffer containing 100 mM NaCl, 50 mM Tris-HCl (pH 7.5), 6 mM MgCl $_2$, 0.5 mM CaCl $_2$, 0.5 mM dithiothreitol, 0.05% Nonidet P-40. After 30 min incubation at 30°C , reactions were terminated by adding 100 μl GREEN reagent. A $_{620}$ was

measured using a microtiter plate reader. After background subtraction from each sample, the activity of calcineurin was determined as the difference between total phosphatase activity and the phosphatase activity in the EGTA containing buffer.

Statistics

All statistical analyses were performed with Prism 5 software (GraphPad Software). The normal distribution of the data was assessed using D'Agostino-Pearson omnibus test. Accordingly, parametric or non-parametric test was applied (as indicated in each experiment). In all graphs numbers within bars depict the numbers of analyzed individual visual fields (= cells) or independent wells/samples obtained from at least two different cell culture preparations. All data are always normalized to the mean of the control group and expressed as mean \pm SEM. The level of statistical significance is indicated as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ in all the graphs.

RESULTS

Th Modulates Recycling of SVs at Excitatory and Inhibitory Synapses

In order to investigate the role of endogenously secreted A β in the regulation of neurotransmitter release, we applied Th to the cultured rat cortical neurons at DIV18–21. Th is an inhibitor of metalloprotease neprilysin, which mediates the rate-limiting step in A β degradation (Iwata et al., 2001; Abramov et al., 2009). Treatment with Th for 1 h led to 1.6 and 1.2 fold elevation in the extracellular concentrations of A β_{42} and A β_{40} , respectively compared to untreated control, as assessed by sandwich ELISA of freshly collected neuronal growth media (**Figure 1A**; [A β_{42}]: control: 57 ± 11 pM vs. Th: 94 ± 15 pM, **Figure 1B**; [A β_{40}]: control: 130 ± 11 pM vs. Th: 153 ± 11 pM). Low pM concentrations of endogenously produced A β_{42} and A β_{40} peptides measured here as well as their Th-mediated elevation are in line with previous reports and comparable with concentrations obtained *in vivo* (Cirrito et al., 2005; Abramov et al., 2009).

To examine the effect of Th on presynaptic function, we monitored the efficacy of SV recycling at the level of individual synapses using synaptotagmin1 antibody (sytl Ab) uptake (Kraszewski et al., 1995; Lazarevic et al., 2011). This assay utilizes a fluorophore-labeled antibody that recognizes a luminal domain of the SV protein sytl. This epitope becomes accessible for the antibody, added to neuronal media, only after fusion of SVs for neurotransmitter release before they undergo rapid compensatory endocytosis. Network activity-driven sytl Ab uptake was significantly increased in cultures treated with Th compared to untreated control (**Figures 1C,D**; Th $138 \pm 5\%$ of control). Interestingly, application of Th led to an increased network activity-driven recycling of SV in both excitatory and inhibitory synapses as shown by quantification of sytl Ab uptake in puncta immunoreactive for glutamatergic markers homer and vesicular glutamate transporter 1 (VGLUT1) or for a marker of inhibitory synapses, vesicular GABA transporter (VGAT; **Figures 1E,F**; Homer, Th: $146\% \pm 12\%$; VGLUT1, Th: $145 \pm 10\%$; VGAT, Th: $162 \pm 13\%$ of control). Th-mediated

increase of SV recycling in inhibitory synapses was further confirmed by performing uptake assay with an antibody against luminal domain of VGAT (Th: $151 \pm 9\%$ of control).

Extracellular A β_{40} and A β_{42} Exert Hormetic Effect on SV Recycling

To confirm that Th-induced changes in SV recycling rely on the modulation of the extracellular concentration of the endogenously secreted A β peptides, we treated neurons with Th in the presence or absence of 4G8 monoclonal antibody (5 μ g/ml) that specifically binds A β peptides. While incubation with Th alone clearly increased SV recycling, this effect was completely prevented by chelation of extracellular A β using 4G8 antibody (**Figure 2A**; Th: $138 \pm 5\%$; 4G8/Th: $110 \pm 4\%$ of control). Next, we inhibited production of A β , either by blocking β -secretase (BACE inhibitor IV, 0.5 μ M) or γ -secretase (GAMMAinh, L-685, 458, 0.2 μ M). Treatment with these inhibitors for 6 h led to a notable decrease of the sytl Ab uptake, pointing to the role of endogenously released A β in the modulation of basal SV recycling (**Figure 2A**; β inh: $83 \pm 7\%$; γ inh: $73 \pm 5\%$ of control). Moreover, pre-treatment with secretase inhibitors, 5 h prior to the treatment with Th for 1 h, fully blocked the Th-induced increase in SV recycling (β inh/Th: $76 \pm 6\%$; γ inh/Th: $71 \pm 6\%$ of control). Taken together these experiments strongly support involvement of endogenously secreted A β in the Th-induced increase in the sytl Ab uptake.

To strengthen this assumption, we added synthetic A β_{42} and A β_{40} , in physiological concentrations of 200 pM, to the growth media for 1 h, which caused a substantial increase in the sytl Ab uptake (**Figures 2B,C**; A β_{40} : $154 \pm 9\%$; A β_{42} : $157 \pm 9\%$ of control). In contrary, 1-h treatment with peptides from identical preparation at 1 μ M concentration, widely used to induce neurotoxic effects (Park et al., 2013), had an opposite effect and led to a significant decrease in sytl Ab uptake (**Figures 2B,C**; A β_{40} : $69 \pm 4\%$; A β_{42} : $77 \pm 3\%$ of control). None of the treatments affected the density of synapses contacting proximal dendrites assessed as number of puncta immunoreactive for synaptic vesicle protein sph (**Figure 2D**; control: 43 ± 1 ; Th: 43 ± 1 ; 200 pM A β_{40} : 42 ± 2 , A β_{42} : 41 ± 1 ; 1 μ M A β_{40} : 47 ± 3 , A β_{42} : 38 ± 2 synapses). The number of active synapses assessed as sph puncta with over-threshold immunofluorescence for sytl Ab uptake was decreased in cells treated with 1 μ M A β_{42} but unchanged in all other conditions (**Figure 2D**, control: 36 ± 1 ; Th: 39 ± 2 ; 200 pM A β_{40} : 37 ± 3 , A β_{42} : 37 ± 2 ; 1 μ M A β_{40} : 36 ± 3 , A β_{42} : 27 ± 2 active synapses). The percentage of active synapses calculated as proportion of active synapses out of sph positive synapses for each analyzed visual field differed between Th-treated cells and cells treated with 1 μ M A β peptides further indicating that elevated physiological and supraphysiological concentrations of A β_{40} and A β_{42} exert opposite effects on presynaptic function (**Figure 2D**, control: $83 \pm 2\%$; Th: $89 \pm 4\%$; 200 pM A β_{40} : $87 \pm 3\%$, A β_{42} : $89 \pm 3\%$; 1 μ M A β_{40} : $79 \pm 4\%$, A β_{42} : $73 \pm 5\%$). These results are in line with a hormetic effect of A β_{40} and A β_{42} peptides, with low, physiological (high pM) concentration potentiating SV recycling, and high, supraphysiological (low μ M) concentration having a negative impact in the identical experimental readout.




FIGURE 1 | Endogenous A β modulates presynaptic activity at excitatory and inhibitory synapses. **(A,B)** Quantification of A β_{x-42} **(A)** and A β_{x-40} **(B)** levels in the medium from vehicle- and Th-treated cortical cells using sandwich ELISA. Numbers within columns represent the number of individually treated and analyzed wells obtained from two **(A)** and three **(B)** independently prepared cell cultures. **(C)** Representative images of syt1 Ab uptake (magenta) from vehicle- and Th-treated cortical neurons. Synaptophysin (sph, green) was used as a synaptic marker. Regions showed in high magnification are boxed. **(D)** Quantification of normalized fluorescence intensity of network activity-driven syt1 Ab uptake in control and Th-treated neurons along proximal dendrites. **(E)** Representative images of syt1 Ab uptake, from cortical neurons incubated with Th or control solution, at excitatory (homer, green) and inhibitory (VGAT, green) terminals. **(F)** Statistical quantification of the intensity of syt1 Ab uptake in puncta positive for either excitatory (homer, VGLUT) or inhibitory (VGAT) synapses driven by the spontaneous network activity. Numbers within columns show the number of analyzed cells derived from five **(D)** and two **(F)** independent cultures. Values are expressed as the mean \pm SEM. Effect of treatment was compared to untreated control by unpaired two-tailed Student *t* test **(A,B,D,F)** syt1 Ab uptake in VGLUT-positive and VGAT-positive) or Mann Whitney test **(F, syt1 Ab uptake in homer-positive puncta)**; ***p* < 0.01, ****p* < 0.001. Scale bars represent 10 μ m in overview and 5 μ m in insets.

To study durability and reversibility of A β -mediated effect on SV recycling, cells were treated with 200 pM A β_{42} for 1 h prior to the washout and replacement with the conditioned medium. In this experiment, cells monitored 1 h after 200 pM A β_{42} treatment showed a significant increase in syt1 Ab uptake. In contrast, cells assayed 23 h after the A β_{42} washout showed recycling indistinguishable from untreated controls (**Figures 2E,F**; A β_{42} : $141 \pm 5\%$; A β_{42} /washout: $113 \pm 8\%$ of control). This suggests that modulation of A β production and clearance might represent a physiological mechanism, inducing rapid changes in the recycling of SVs.

A β Potentiates Basal Neurotransmission via an Increase in the Recycling Pool of SVs

The observed increase in the network activity-driven SV recycling points to changes in presynaptic properties, but could also reflect elevated network activity in Th- or A β -treated cultures. To test whether this effect relies on a modulation of presynaptic mechanisms, we performed syt1 Ab uptake during pulse-application of 50 mM KCl, which depolarizes neuronal membranes and induces release of all releasable SVs,

i.e., recycling pool (RP) of SVs (Alabi and Tsien, 2012). Depolarization-induced syt1 Ab uptake was upregulated by 40% in Th-treated cultures (**Figures 3A,B**; Th: $142 \pm 8\%$ of control) suggesting the role of the endogenous A β in the regulation of the recycling pool of SV. Accordingly, application of 200 pM A β_{42} exerted similar effect (**Figure 3B**; A β_{42} : $137 \pm 9\%$ of control). Interestingly, high 1 μ M concentration of synthetic A β_{42} significantly reduced the depolarization-induced recycling confirming a hormetic nature of A β -mediated modulation of the presynaptic properties (**Figure 3B**; A β_{42} : $78 \pm 4\%$ of control).

Since syt1 Ab uptake enables only end point measurement of SV recycling and fully relies on the endogenous expression of syt1, we sought to assess whether SV turnover was increased by an alternative method. To this end, we moved to hippocampal neurons, where the potentiating effect of Th, A β_{40} and A β_{42} was originally described by others and reproduced by us in the synapses contacting proximal dendritic segments (**Figures 3C,D**; Th proximal: $139 \pm 13\%$ of proximal control; Th distal: $119 \pm 15\%$ of distal control). We performed live imaging of SV turnover implementing genetically encoded pH-sensitive probe called synaptophysin-pHluorin (sypHy; Burrone et al., 2006; Granseth et al., 2006; Rose et al., 2013) expressed in





FIGURE 2 | Extracellular A β_{40} and A β_{42} have a homeostatic effect on presynaptic vesicle recycling. **(A)** Quantification of activity-driven syt1 Ab uptake from Th-treated vs. control cells in the presence or absence of A β antibody (4G8) and inhibitors of enzymes involved in amyloid precursor protein (APP) processing (β -secretase and γ -secretase inhibitors). Numbers within bars display the number of analyzed visual fields pooled from five (Th), three (β inh; γ inh) and two (4G8) independent cell culture preparations. **(B)** Representative images of cultured cortical neurons treated either with pM or μ M concentration of A β_{40} , A β_{42} or control solution and stained with syt1 Ab (magenta) and sph (green). **(C)** Quantification of **(B)**. Numbers indicate the number of individual cells used for analysis obtained from four (A β_{42}) and two (A β_{40}) culture preparation. **(D)** Quantification of density of sph-positive synapses (white bars), active synapses with syt1 Ab uptake (gray bars) and quantification of percentage of active synapses (black bars) in cultures treated with vehicle, Th and 200 pM and 1 μ M A β_{40} and A β_{42} . Numbers in columns indicate number of analyzed cells coming from at least two independent cell preparations. **(E)** Representative images of cortical neurons treated with control or 200 pM A β_{42} as well as cells assayed 23 h after A β_{42} washout for network activity-driven syt1 Ab uptake. **(F)** Quantification of the experiment in **(E)**. Numbers within columns show the number of the cells used for analysis from at least two independent cell culture preparations. In all graphs values are expressed as the mean \pm SEM. The density of synapses were analyzed using one-way ANOVA with Bonferroni *post hoc* test, all other data were analyzed using Kruskal-Wallis test followed by Dunn's multiple comparison test, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Scale bar, 5 μ m.



cultured hippocampal neurons using lentiviral vectors. SytHy fluorescence is quenched at low, acidic pH found inside the SVs, but increases upon SV fusion and exposure of their lumen to the neutral pH of the extracellular media. The imaging was performed in the presence of bafilomycin, an inhibitor of the vesicular proton pump that prevents reacidification of SVs upon their compensatory endocytosis. To visualize readily releasable pool (RRP), which corresponds to the morphologically docked SVs, we delivered 40 AP at 20 Hz (Burrone et al., 2006). After 2 min pause to allow for the recovery of RRP, 900 AP at 20 Hz were delivered to release all releasable SVs (RP). SV refractory to stimulation (resting pool, RtP) were

visualized by the alkalization of the SVs lumen using NH $_4$ Cl pulse, which enables accurate assessment of RP relative to all SVs present at the individual synapse (Burrone et al., 2006). In neurons treated with A β_{42} no changes in RRP, but a significant increase of RP (**Figures 3E,F**; A β_{42} : 112 \pm 1% of control) and correspondingly a decrease in RtP were observed. These experiments ultimately demonstrate that A β modifies presynaptic function by the regulation of the turnover of presynaptic SV.

Role of Calcium Signaling via α 7nAChR in the A β -Mediated Potentiation of SV Recycling

Previous studies proposed that, at picomolar concentrations, A β could bind and activate presynaptic α 7nAChR (Wang et al., 2000b; Dougherty et al., 2003). Thus, we sought to determine the role of these receptors in the A β -mediated regulation of the presynaptic neurotransmitter release. Treatment with α -bungarotoxin (BgTx, 50 nM), a specific blocker of α 7nAChRs, for 90 min did not significantly affect syt1 Ab uptake in untreated cells. However, the same treatment completely prevented increase in syt1 Ab uptake induced by Th- or 200 pM A β_{42} application for 1 h, revealing a critical role of α 7nAChR in the A β_{42} -mediated regulation of SV cycling (**Figures 4A,B**; BgTx: 91 \pm 5%; Th: 151 \pm 11%; BgTx/Th: 109 \pm 8%; A β_{42} : 127 \pm 6%; BgTx/A β_{42} : 101 \pm 9% of control). Interestingly, BgTx application was unable to completely block decreased syt1 Ab uptake mediated by supraphysiological concentration (1 μ M) of A β_{42} (**Figures 4A,B**; BgTx: 89 \pm 6%; A β_{42} : 68 \pm 5%; BgTx/A β_{42} : 81 \pm 4%). This indicates that the effect mediated by higher concentrations of the peptide may involve not only α 7nAChRs-dependent signaling, but also other types of either pre- or postsynaptic receptors (Lauren et al., 2009; Nikolaev et al., 2009). Next, we wondered whether activation of α 7nAChR is sufficient to mimic Th-induced upregulation of syt1 Ab uptake. To this end, we applied choline (Ch, 500 μ M), an agonist of α 7nAChR (Alkondon et al., 1997), 20 min prior to Th or control treatment. Interestingly, choline had no effect on network activity-driven syt1 Ab uptake in control cells, however it completely blocked Th-induced presynaptic activity when applied together with Th (**Figure 4C**; Ch: 110 \pm 9%; Th: 149 \pm 9%; Ch/Th: 91 \pm 5% of control). This might be due to the well-known, fast desensitization of AChRs upon choline binding. To confirm this, we tested the impact of PNU120596, an allosteric modulator of α 7nAChR, which has been shown to increase mean opening time of these receptors and thereby interfere with channel desensitization (Hurst et al., 2005). PNU120596 (3 μ M, 1 h), had no effect on syt1 Ab uptake (**Figure 4C**; PNU: 114 \pm 6% of control). Co-application of PNU120596 with choline fully mimicked Th-induced response (**Figure 4C**; PNU/Ch: 155 \pm 11% of control) but PNU120596 was unable to further potentiate the effect of Th (**Figure 4C**; PNU/Th: 145 \pm 15% of control), suggesting that endogenous A β might influence both channel opening and stabilization. Importantly, the Th-mediated increase in the depolarization-induced syt1 Ab





FIGURE 4 | Picomolar A β acts via α 7nACh receptors. **(A)** Representative images of syt1 Ab uptake from Th-, A β ₄₂ 200 pM or A β ₄₂ 1 μ M-treated neurons pre-incubated with BgTx to investigate the effect of α 7nAChR. **(B)** Quantification of syt1 Ab uptake from **(A)**. Numbers within columns indicate numbers of analyzed cells from two (Th; A β ₄₂ 200 pM) or three (A β ₄₂ 1 μ M) cell culture preparations. **(C)** Statistical analysis of syt1 Ab uptake from cortical neurons treated with control, Th or/and ortho- and allosteric modulators of α 7nACh receptors. **(D)** Quantification of depolarization-induced syt1 Ab uptake after control, Th and/or BgTx application. The numbers within bars **(C,D)** indicate the number of analyzed cells obtained from two separate cell culture preparations. In all graphs the values are expressed as the mean \pm SEM. Statistical significance was evaluated by Kruskal-Wallis test followed by Dunn's multiple comparison test **(B,C)** or one-way ANOVA with Bonferroni *post hoc* test **(D)**; * p < 0.05, ** p < 0.01, *** p < 0.001. Scale bar 5 μ m.

uptake was completely precluded by preincubation of cells, for 30 min, with BgTx before Th application for further 1 h (**Figure 4D**; Th: $140 \pm 7\%$; BgTx: $108 \pm 8\%$; BgTx/Th:

$100 \pm 7\%$ of control). Our data reveal that the A β -induced potentiation of presynaptic function relies on the modulation of α 7nAChRs.



One of the most prominent features of α 7nACh receptors is their high Ca^{2+} conductance, which substantially contributes to synaptic Ca^{2+} signaling. To investigate whether influx of extracellular Ca^{2+} plays a role in the potentiation of SV recycling mediated by Th, we incubated both control and Th-treated cells in Ca^{2+} -free medium for 1 h and subsequently performed live staining with syt1 Ab in usual imaging media containing 2 mM Ca^{2+} . Incubation of cells in the Ca^{2+} -free medium produced no significant effect on the network activity-driven syt1 Ab uptake (Figure 5A; Ctrl/no Ca^{2+} : $99 \pm 8\%$ of control) nor on the size of RP assessed by syt1 Ab uptake upon pulse application of 50 mM KCl (Figure 5B; Ctrl/no Ca^{2+} : $89 \pm 4\%$ of control). However, the impact of Th treatment on both network activity-driven and depolarization-induced

syt1 Ab uptake was fully abolished in the Ca^{2+} -free medium (Figures 5A,B; network activity driven syt1 Ab uptake: Th: $136 \pm 8\%$; Th/no Ca^{2+} : $96 \pm 7\%$ of control; syt1 Ab uptake upon KCl depolarization: Th: $139 \pm 8\%$; Th/no Ca^{2+} : $89 \pm 4\%$ of control).

α 7nAChRs are also essentially involved in the activation of the calcium-induced calcium release pathway, which governs calcium efflux from cellular internal stores via stimulation of ryanodine receptors (RyR; Sharma and Vijayaraghavan, 2001; Dajas-Bailador et al., 2002). To test the involvement of this signaling in Th-mediated presynaptic strengthening, we applied a RyR blocker, TMB-8 (100 μM , 1 h), to the control and Th-treated cells and quantified their network activity-driven syt1 Ab uptake. The TMB-8-mediated block of Ca^{2+} release from the intracellular stores efficiently impeded Th-induced effect on the SV recycling (Figure 5C; Th: $153 \pm 11\%$; TMB-8/Th: $76 \pm 5\%$; TMB-8: $107 \pm 9\%$ of control). Furthermore, application of cell-permeable calcium chelator, BAPTA-AM (10 μM , 1 h), also prevented the increase in SV recycling in Th-treated cells (Figure 5D; Th: $133 \pm 8\%$; BAPTA-AM/Th: $89 \pm 7\%$; BAPTA-AM: $76 \pm 5\%$ of control), suggesting contribution of calcium-dependent pathways in A β -driven presynaptic strengthening.

Involvement of CDK5/Calcineurin Balance in the A β -Mediated Regulation of SV Pools

In the recent years, cyclin dependent kinase 5 (CDK5) and phosphatase calcineurin emerged as two major players controlling the efficacy of neurotransmitter release by modulation of the size of RP of SVs (Kim and Ryan, 2010, 2013; Marra et al., 2012). Moreover, CDK5 and calcineurin are well described targets of calcium signaling and were already previously linked to downstream signaling via α 7nAChR (Patrick et al., 1999; Lee et al., 2000; Stevens et al., 2003). To address potential contribution of CDK5 and calcineurin in A β -mediated alternations in SV recycling, we pharmacologically blocked these enzymes and carried out syt1 Ab uptake under high KCl conditions in which all recycling vesicles undergo exocytosis and are labeled. Application of Roscovitine (1 h, 50 μM), a potent CDK5 inhibitor (Meijer et al., 1997) led to a considerable increase of syt1 Ab uptake, yet combined treatment (Th/Roscovitine) did not exert any further effect (Figure 6A; Th: $136 \pm 6\%$; Roscovitine: $161 \pm 8\%$; Th/Roscovitine: $132 \pm 7\%$ of control). In contrast, inhibition of calcineurin by FK506 (1 μM , 1 h) induced a significant reduction in the depolarization-induced syt1 Ab uptake in both, control and Th-treated cells (Figure 6B; Th: $129 \pm 7\%$; FK506: $86 \pm 6\%$; Th/FK506: $105 \pm 5\%$ of control). These results are in line with possible involvement of CDK5/calcineurin signaling in A β -driven regulation of SV recycling pool. To further explore this hypothesis we performed CDK5 and calcineurin activity assay. Analysis of kinase activity confirmed that cells treated with Th or 200 pM A β_{42} , show significant reduction in CDK5 activity (Figure 6C; Th: $80 \pm 1\%$; A β_{42} : $85 \pm 1\%$ of control), with no change in the total protein levels (Figure 6D). Furthermore, Th-induced effect was completely prevented when α 7nAChR were blocked (Figure 6C; BgTx/Th:




FIGURE 6 | CDK5 and calcineurin are involved in A β -driven presynaptic plasticity. **(A)** Quantification of depolarization-induced syt1 Ab uptake from cortical neuronal cultures treated with control, Th and/or roscovitine (Rosc) to evaluate the impact of CDK5. **(B)** Statistical analysis of depolarization-induced syt1 Ab uptake from cortical cells incubated with vehicle, Th and/or FK506 to investigate the relevance of calcineurin signaling. Numbers within columns in **(A,B)** represent the number of analyzed cells obtained from three independent cell culture preparations. **(C)** CDK5 activity assay performed from cortical cultures under different conditions. **(D)** Representative Western blot as well as quantification of the total CDK5 protein levels from vehicle- or Th-treated cultures. Molecular weight is indicated in kDa. The numbers represent the number of samples from two independent primary culture preparations. **(E)** Calcineurin activity assay conducted from cortical cells incubated with vehicle or Th. Numbers within columns in **(C,E)** denote the number of independently treated and analyzed wells in a multi-well dishes obtained from three **(C)** or four **(E)** different cell culture preparations. In all graphs, values are expressed as the mean \pm SEM. Statistical significance was assessed by one-way ANOVA with Bonferroni *post hoc* test **(A–C)** or Student **(D)** and Mann Whitney *t* test **(E)**; ** $p < 0.01$, *** $p < 0.001$.

101 \pm 2% of control). On the other hand, a phosphatase activity assay revealed significantly higher calcineurin activity in the neurons treated with Th (**Figure 6E**; Th: 179 \pm 14% of control), corroborating that balancing the activity of these enzymes has an important role in A β -driven regulation of the recycling vesicles. Altogether, these results support a view that Th-mediated regulation of SV turnover involves modulation of CDK5/calcineurin phosphohomeostasis downstream of α 7nAChR receptors.

DISCUSSION

Despite the central role of A β peptide in the etiopathogenesis of AD, its physiological function in the healthy brain is still poorly understood. During the past decades, A β emerged as a vital factor that regulates neurotransmission and studies exploring effects of physiological i.e., low-intermediate picomolar concentrations of A β suggested presynaptic locus of its action. In this work, we directly tested the role of Th, A β ₄₀ and A β ₄₂ species in the

regulation of neurotransmitter release from the presynapse and examined the underlying cellular signaling.

Th and Endogenous A β_{40} and A β_{42} Converge on Regulation of SV Recycling

Using quantification of syt1 Ab uptake at levels of individual synapses, we confirmed previously reported effects of Th on SV recycling in cultured rat cortical and hippocampal neurons. The authors of the original publication argued that Th effect is based on the elevation of extracellular concentrations of endogenously produced A β species and provided numerous, yet indirect, evidence for their claim (Abramov et al., 2009). In a following study, presynaptic strengthening was induced in the same neuronal cultures by human A β_{40} , which, however, differs from its murine ortholog in its N-terminal sequence (Fogel et al., 2014). Thus, the presynaptic effect of endogenous A β_{40} and A β_{42} has never been explicitly demonstrated and compared. Our experiments extend previous findings and show that Th and 200 pM A β_{40} or A β_{42} , in parallel experiments, exert the same effect, namely an increase in the turnover of SVs. All treatments could be efficiently blocked by inhibition of α 7-nAChRs, which strongly speaks for their common mechanism of action. Interestingly, we observed smaller relative increase in the size of the recycling pool of SVs compared to the effect seen in the depolarization-driven syt1 Ab uptake assays, which might reflect differences in vesicle origin released by the chemical and electrical stimuli. Accelerated compensatory endocytosis of released SVs could also contribute to this effect and should be tested in future studies. In contrast to the previous studies that argued against any impact of physiological A β on the inhibitory synapses, we observed an A β treatment-induced increase in the network activity-driven recycling of SVs in the inhibitory synapses (Abramov et al., 2009). However, we cannot exclude that effect shown here (Figure 1F) simply reflects an increase in the overall network activity upon Th treatment and not direct regulation of release at inhibitory synapses.

Calcium Influx via α 7nAChR is Required for A β -Mediated Increase in SV Recycling

Involvement of α 7nAChR in the modulation of neurotransmission by A β is controversial. The human and rodent A β_{42} -induced memory enhancement and LTP increase were absent in the knock out for the α 7 subunit of nAChR and A β_{42} -induced LTP was also sensitive to the antagonists of α 7nAChR BgTx and mecamylamine (Puzzo et al., 2008, 2011). In contrast, the presynaptic strengthening shown by imaging of activity-induced styryl dye destaining upon treatment with Th and human A β_{40} was not sensitive to pharmacological block of these receptors (Fogel et al., 2014). In the experiments described here, the effect of Th and picomolar concentrations of A β_{42} on network activity- and depolarization-driven SV recycling was fully prevented upon pretreatment with α 7nAChR competitive antagonist-BgTx. In line with a requirement of the α 7nAChR-mediated calcium influx for the Th-induced increase in SV recycling the effect of Th was absent upon depletion of extracellular and/or intracellular calcium and upon interference

with calcium-induced calcium release from cellular internal stores. The effect of Th was also lost in the cells pretreated with choline, an agonist of α 7nAChR that at high concentrations induces a rapid desensitization of the channel. Interestingly, a co-application of desensitizing concentration of choline and the allosteric modulator of α 7nAChR PNU120596, known to increase opening time of agonist-bound receptor and decrease the receptor desensitization was comparable to the Th-mediated increase in SV recycling. Co-application of PNU120596 and Th did not further potentiate SV recycling suggesting that Th might affect receptor desensitization kinetics similarly as PNU120596. However, it is still unclear, whether A β_{40} and A β_{42} act as agonists or modulators of α 7nAChR and what are the cofactors required for their action. While the impact of 200 pM A β_{40} and A β_{42} was hindered by pharmacological interference with α 7nAChR, a blockage of these receptors did not fully prevent the effect of 1 μ M A β_{42} . This might imply that at higher concentrations A β_{42} acts via different cell receptors. Nevertheless, displacement measurements of A β_{42} and BgTx on radiolabeled α 7nAChR suggested a competition of the two compounds on the same binding site with A β_{42} having a 4000-fold higher affinity as compared to BgTx (Wang et al., 2000b). Therefore, our result might be also explained by an incomplete blockage of α 7nAChR with BgTx in the presence of 1 μ M A β_{42} .

Endogenous A β Modulates Recycling SV via Modulation of CDK5 and Calcineurin Activity

Quantification of depolarization-induced syt1 Ab uptake and syPhy imaging in living neurons shown here strongly implies that Th, A β_{40} and A β_{42} control neurotransmitter release via regulation of the recycling of SVs. Recently, CDK5 and calcineurin were suggested to govern the recycling of SVs by setting a balance in the local phospho- and dephosphorylation (Kim and Ryan, 2010, 2013; Marra et al., 2012). We have shown that application of Th or A β_{42} leads to a rapid decrease of CDK5 activity in the cell lysates. This decrease in CDK5 activity required normal α 7nAChR signaling substantiating their function upstream of CDK5 in the regulation of SV recycling by A β . Pharmacological inhibition of CDK5 by roscovitine mimicked Th-induced increase in the depolarization-induced syt1 Ab uptake and co-application of Th and roscovitine had no additive effect, suggesting that they share common pathways in regulation of the SV turnover. Th application increased the calcineurin activity and a pharmacological inhibition of calcineurin activity interfered with the Th-induced increase in the depolarization-induced recycling of SVs. At present, we can only speculate about signaling pathways connecting calcium influx via α 7nAChR and regulation of calcineurin and CDK5 activity. Previously reported activation of calcineurin by calcium influx through α 7nAChRs is compatible with our observations (Stevens et al., 2003). Cleavage of p35, an activator of CDK5, was observed upon application of micromolar A β_{42} and resulted in a formation of CDK5/p25 hyperactive complex (Patrick et al., 1999; Lee et al., 2000). This scenario contradicts our results that demonstrate calcium influx-induced decrease of CDK5 activity. It is possible that a slight elevation of intracellular

calcium upon A β -mediated activation of $\alpha 7$ nAChR has different consequences for CDK5 regulation than a massive calcium influx induced by a prolonged application of micromolar A β_{42} .

Taken together, our data support the function of endogenous A β species in the regulation of neurotransmitter release. The described modulation of presynaptic function by A β was rapid and reversible. Moreover, a depletion of endogenous A β upon interference with its production and application of elevated A β concentrations led to a decrease in the synaptic strength, which is in accordance with previously proposed hormetic regulation of neurotransmission by endogenous A β (Puzzo et al., 2008; Abramov et al., 2009). Thus, our results corroborate on function of endogenous A β as a physiological regulator of presynaptic efficacy. Compellingly, the fast modulation of release by A β might act in tuning of synaptic strength at the level of individual synapses in processes of synaptic plasticity. The observation that intracellular signaling cascades, shown here to be involved in the physiological regulation of SV recycling by A β , are also implicated in AD pathogenesis supports the hypothesis that failure of the physiological function of A β in the tuning of SV recycling could impair synaptic homeostasis and initiate synaptic dysfunction leading to cognitive decline and neurodegeneration in AD.

AUTHOR CONTRIBUTIONS

VL and AF conceptualized the study and curated the data. VL, SF, MA-A and DA performed investigations, formal analysis,

methodology development and validation. CM-V, DI, MAC provided methodologies. AF and EDG provided infrastructure and resources. VL, SF and MA-A executed visualization of data. AF supervised the study. VL, SF and AF have written original draft. All authors reviewed and edited the final manuscript.

FUNDING

This study was supported by grants from following agencies: Deutsche Forschungsgemeinschaft (FE1335/1 and SFB779/A06), People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ NPlast under REA grant agreement no. [289581], Federal State of Saxony-Anhalt and the European Regional Development Fund (ERDF) (CBBS, FKZ: ZS/2016/04/78120), Leibniz Association (LGS Synaptogenetics, SAW 2013-15 and SAW 2014-2016). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

ACKNOWLEDGMENTS

We thank I. Herbert, K. Hartung, B. Kracht and S. Müller for help with primary neuronal cultures, T. Oertner for the syHy construct, N. Ziv for providing the OpenView software, P. Lewczuk for recommendations on ELISA measurements and R. Frischknecht and all lab members for conceptual discussions.

REFERENCES

- Abramov, E., Dolev, I., Fogel, H., Cicciotosto, G. D., Ruff, E., and Slutsky, I. (2009). Amyloid- β as a positive endogenous regulator of release probability at hippocampal synapses. *Nat. Neurosci.* 12, 1567–1576. doi: 10.1038/nn.2433
- Alabi, A. A., and Tsien, R. W. (2012). Synaptic vesicle pools and dynamics. *Cold Spring Harb. Perspect. Biol.* 4:a013680. doi: 10.1101/cshperspect.a013680
- Alkondon, M., Pereira, E. F., Cortes, W. S., Maelicke, A., and Albuquerque, E. X. (1997). Choline is a selective agonist of $\alpha 7$ nicotinic acetylcholine receptors in the rat brain neurons. *Eur. J. Neurosci.* 9, 2734–2742. doi: 10.1111/j.1460-9568.1997.tb01702.x
- Banker, G. A. (1980). Trophic interactions between astroglial cells and hippocampal neurons in culture. *Science* 209, 809–810. doi: 10.1126/science.7403847
- Burrone, J., Li, Z., and Murthy, V. N. (2006). Studying vesicle cycling in presynaptic terminals using the genetically encoded probe synaptopHluorin. *Nat. Protoc.* 1, 2970–2978. doi: 10.1038/nprot.2006.449
- Chapman, P. F., White, G. L., Jones, M. W., Cooper-Blacketer, D., Marshall, V. J., Irizarry, M., et al. (1999). Impaired synaptic plasticity and learning in aged amyloid precursor protein transgenic mice. *Nat. Neurosci.* 2, 271–276. doi: 10.1038/6374
- Cirrito, J. R., May, P. C., O'dell, M. A., Taylor, J. W., Parsadanian, M., Cramer, J. W., et al. (2003). *In vivo* assessment of brain interstitial fluid with microdialysis reveals plaque-associated changes in amyloid- β metabolism and half-life. *J. Neurosci.* 23, 8844–8853.
- Cirrito, J. R., Yamada, K. A., Finn, M. B., Sloviter, R. S., Bales, K. R., May, P. C., et al. (2005). Synaptic activity regulates interstitial fluid amyloid- β levels *in vivo*. *Neuron* 48, 913–922. doi: 10.1016/j.neuron.2005.10.028
- Crews, L., Patrick, C., Adame, A., Rockenstein, E., and Masliah, E. (2011). Modulation of aberrant CDK5 signaling rescues impaired neurogenesis in models of Alzheimer's disease. *Cell Death Dis.* 2:e120. doi: 10.1038/cddis.2011.2
- Dajas-Bailador, F. A., Mogg, A. J., and Wonnacott, S. (2002). Intracellular Ca^{2+} signals evoked by stimulation of nicotinic acetylcholine receptors in SH-SY5Y cells: contribution of voltage-operated Ca^{2+} channels and Ca^{2+} stores. *J. Neurochem.* 81, 606–614. doi: 10.1046/j.1471-4159.2002.00846.x
- Dougherty, J. J., Wu, J., and Nichols, R. A. (2003). β -amyloid regulation of presynaptic nicotinic receptors in rat hippocampus and neocortex. *J. Neurosci.* 23, 6740–6747.
- Fejtova, A., Davydova, D., Bischof, F., Lazarevic, V., Altmann, W. D., Romorini, S., et al. (2009). Dynein light chain regulates axonal trafficking and synaptic levels of Bassoon. *J. Cell Biol.* 185, 341–355. doi: 10.1083/jcb.200807155
- Fogel, H., Frere, S., Segev, O., Bharill, S., Shapira, I., Gazit, N., et al. (2014). APP homodimers transduce an amyloid- β -mediated increase in release probability at excitatory synapses. *Cell Rep.* 7, 1560–1576. doi: 10.1016/j.celrep.2014.04.024
- Frischknecht, R., Fejtova, A., Viesti, M., Stephan, A., and Sonderegger, P. (2008). Activity-induced synaptic capture and exocytosis of the neuronal serine protease neurotrypsin. *J. Neurosci.* 28, 1568–1579. doi: 10.1523/jneurosci.3398-07.2008
- Garcia-Osta, A., and Alberini, C. M. (2009). Amyloid β mediates memory formation. *Learn. Mem.* 16, 267–272. doi: 10.1101/lm.1310209
- Glennner, G. G., and Wong, C. W. (1984). Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem. Biophys. Res. Commun.* 120, 885–890. doi: 10.1016/s0006-291x(84)80190-4
- Goate, A., Chartier-Harlin, M. C., Mullan, M., Brown, J., Crawford, F., Fidani, L., et al. (1991). Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349, 704–706. doi: 10.1038/349704a0

- Graneth, B., Odermatt, B., Royle, S. J., and Lagnado, L. (2006). Clathrin-mediated endocytosis is the dominant mechanism of vesicle retrieval at hippocampal synapses. *Neuron* 51, 773–786. doi: 10.1016/j.neuron.2006.08.029
- Hsia, A. Y., Masliah, E., McConlogue, L., Yu, G. Q., Tatsuno, G., Hu, K., et al. (1999). Plaque-independent disruption of neural circuits in Alzheimer's disease mouse models. *Proc. Natl. Acad. Sci. U S A* 96, 3228–3233. doi: 10.1073/pnas.96.6.3228
- Hsieh, H., Boehm, J., Sato, C., Iwatsubo, T., Tomita, T., Sisodia, S., et al. (2006). AMPAR removal underlies A β -induced synaptic depression and dendritic spine loss. *Neuron* 52, 831–843. doi: 10.1016/j.neuron.2006.10.035
- Hurst, R. S., Hajos, M., Raggenbass, M., Wall, T. M., Higdon, N. R., Lawson, J. A., et al. (2005). A novel positive allosteric modulator of the α 7 neuronal nicotinic acetylcholine receptor: *in vitro* and *in vivo* characterization. *J. Neurosci.* 25, 4396–4405. doi: 10.1523/jneurosci.5269-04.2005
- Iwata, N., Tsubuki, S., Takaki, Y., Shirotani, K., Lu, B., Gerard, N. P., et al. (2001). Metabolic regulation of brain A β by neprilysin. *Science* 292, 1550–1552. doi: 10.1126/science.1059946
- Kamenetz, F., Tomita, T., Hsieh, H., Seabrook, G., Borchelt, D., Iwatsubo, T., et al. (2003). APP processing and synaptic function. *Neuron* 37, 925–937. doi: 10.1016/s0896-6273(03)00124-7
- Kim, S. H., and Ryan, T. A. (2010). CDK5 serves as a major control point in neurotransmitter release. *Neuron* 67, 797–809. doi: 10.1016/j.neuron.2010.08.003
- Kim, S. H., and Ryan, T. A. (2013). Balance of calcineurin A α and CDK5 activities sets release probability at nerve terminals. *J. Neurosci.* 33, 8937–8950. doi: 10.1523/jneurosci.4288-12.2013
- Kraszewski, K., Mundigl, O., Daniell, L., Verderio, C., Matteoli, M., and De Camilli, P. (1995). Synaptic vesicle dynamics in living cultured hippocampal neurons visualized with CY3-conjugated antibodies directed against the lumenal domain of synaptotagmin. *J. Neurosci.* 15, 4328–4342.
- Lauren, J., Gimbel, D. A., Nygaard, H. B., Gilbert, J. W., and Strittmatter, S. M. (2009). Cellular prion protein mediates impairment of synaptic plasticity by amyloid- β oligomers. *Nature* 457, 1128–1132. doi: 10.1038/nature07761
- Lazarevic, V., Schöne, C., Heine, M., Gundelfinger, E. D., and Fejtova, A. (2011). Extensive remodeling of the presynaptic cytomatrix upon homeostatic adaptation to network activity silencing. *J. Neurosci.* 31, 10189–10200. doi: 10.1523/jneurosci.2088-11.2011
- Lee, M. S., Kwon, Y. T., Li, M., Peng, J., Friedlander, R. M., and Tsai, L. H. (2000). Neurotoxicity induces cleavage of p35 to p25 by calpain. *Nature* 405, 360–364. doi: 10.1038/35012636
- Lois, C., Hong, E. J., Pease, S., Brown, E. J., and Baltimore, D. (2002). Germline transmission and tissue-specific expression of transgenes delivered by lentiviral vectors. *Science* 295, 868–872. doi: 10.1126/science.1067081
- Marra, V., Burden, J. J., Thorpe, J. R., Smith, I. T., Smith, S. L., Häusser, M., et al. (2012). A preferentially segregated recycling vesicle pool of limited size supports neurotransmission in native central synapses. *Neuron* 76, 579–589. doi: 10.1016/j.neuron.2012.08.042
- Meijer, L., Borgne, A., Mulner, O., Chong, J. P., Blow, J. J., Inagaki, N., et al. (1997). Biochemical and cellular effects of roscovitine, a potent and selective inhibitor of the cyclin-dependent kinases cdc2, cdk2 and cdk5. *Eur. J. Biochem.* 243, 527–536. doi: 10.1111/j.1432-1033.1997.t01-2-00527.x
- Mucke, L., Masliah, E., Yu, G. Q., Mallory, M., Rockenstein, E. M., Tatsuno, G., et al. (2000). High-level neuronal expression of A β _{1–42} in wild-type human amyloid protein precursor transgenic mice: synaptotoxicity without plaque formation. *J. Neurosci.* 20, 4050–4058.
- Nikolaev, A., McLaughlin, T., O'Leary, D. D., and Tessier-Lavigne, M. (2009). APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature* 457, 981–989. doi: 10.1038/nature07767
- Palop, J. J., and Mucke, L. (2010). Amyloid- β -induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. *Nat. Neurosci.* 13, 812–818. doi: 10.1038/nn.2583
- Park, J., Jang, M., and Chang, S. (2013). Deleterious effects of soluble amyloid- β oligomers on multiple steps of synaptic vesicle trafficking. *Neurobiol. Dis.* 55, 129–139. doi: 10.1016/j.nbd.2013.03.004
- Patrick, G. N., Zukerberg, L., Nikolic, M., de la Monte, S., Dikkes, P., and Tsai, L. H. (1999). Conversion of p35 to p25 deregulates Cdk5 activity and promotes neurodegeneration. *Nature* 402, 615–622. doi: 10.1038/45159
- Puzzo, D., Privitera, L., Fa', M., Staniszewski, A., Hashimoto, G., Aziz, F., et al. (2011). Endogenous amyloid- β is necessary for hippocampal synaptic plasticity and memory. *Ann. Neurol.* 69, 819–830. doi: 10.1002/ana.22313
- Puzzo, D., Privitera, L., Leznik, E., Fa', M., Staniszewski, A., Palmeri, A., et al. (2008). Picomolar amyloid- β positively modulates synaptic plasticity and memory in hippocampus. *J. Neurosci.* 28, 14537–14545. doi: 10.1523/JNEUROSCI.2692-08.2008
- Rose, T., Schoenberger, P., Jezek, K., and Oertner, T. G. (2013). Developmental refinement of vesicle cycling at Schaffer collateral synapses. *Neuron* 77, 1109–1121. doi: 10.1016/j.neuron.2013.01.021
- Sharma, G., and Vijayaraghavan, S. (2001). Nicotinic cholinergic signaling in hippocampal astrocytes involves calcium-induced calcium release from intracellular stores. *Proc. Natl. Acad. Sci. U S A* 98, 4148–4153. doi: 10.1073/pnas.071540198
- Stevens, T. R., Krueger, S. R., Fitzsimonds, R. M., and Picciotto, M. R. (2003). Neuroprotection by nicotine in mouse primary cortical cultures involves activation of calcineurin and L-type calcium channel inactivation. *J. Neurosci.* 23, 10093–10099.
- Terry, R. D., Masliah, E., Salmon, D. P., Butters, N., Deteresa, R., Hill, R., et al. (1991). Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. *Ann. Neurol.* 30, 572–580. doi: 10.1002/ana.410300410
- Tsuriel, S., Geva, R., Zamorano, P., Dresbach, T., Boeckers, T., Gundelfinger, E. D., et al. (2006). Local sharing as a predominant determinant of synaptic matrix molecular dynamics. *PLoS Biol.* 4:e271. doi: 10.1371/journal.pbio.0040271
- Walsh, D. M., Klyubin, I., Fadeeva, J. V., Cullen, W. K., Anwyl, R., Wolfe, M. S., et al. (2002). Naturally secreted oligomers of amyloid β protein potently inhibit hippocampal long-term potentiation *in vivo*. *Nature* 416, 535–539. doi: 10.1038/416535a
- Wang, H. Y., Lee, D. H., D'Andrea, M. R., Peterson, P. A., Shank, R. P., and Reitz, A. B. (2000a). β -Amyloid_{1–42} binds to α 7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J. Biol. Chem.* 275, 5626–5632. doi: 10.1074/jbc.275.8.5626
- Wang, H. Y., Lee, D. H., Davis, C. B., and Shank, R. P. (2000b). Amyloid peptide A β _{1–42} binds selectively and with picomolar affinity to α 7 nicotinic acetylcholine receptors. *J. Neurochem.* 75, 1155–1161. doi: 10.1046/j.1471-4159.2000.0751155.x

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Lazarevic, Fieńko, Andres-Alonso, Anni, Ivanova, Montenegro-Venegas, Gundelfinger, Cousin and Fejtova. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.